

Research Article

Association between Academic Burnout and Physiological Stress Indicators in Medical Students

Samia Sarwar^{1*}, Sobia Nawaz², Iram Qamar³, Syed Hussain Raza⁴, Junaid Iqbal⁵, Sehrish Liaqat⁶

^{1*}Professor, Department of Physiology, Rawalpindi Medical University Rawalpindi, Pakistan.

²Assistant Professor, Department of Medical Education, Continental Medical College Lahore, Pakistan.

³Professor, Department of Physiology, Queens Medical College Kasur, Pakistan.

⁴Assistant Professor, Department of Medical Education, Avicenna Medical College Lahore, Pakistan.

⁵Associate Professor, Department of Physiology, Sahara Medical College Narowal, Pakistan.

⁶Senior Lecturer, Department of Community Medicine, Sahara Medical College Narowal, Pakistan.

Corresponding Author: Samia Sarwar

Email: drsamsarwar@yahoo.com

Received: 05.10.25, Revised: 07.11.25, Accepted: 11.01.26

ABSTRACT

Background: Academic burnout is increasingly recognized as a significant concern among medical students due to prolonged academic demands, high expectations, and psychological pressure. Burnout is commonly characterized by emotional exhaustion, depersonalization, and reduced personal accomplishment. While its psychological impact has been widely studied, its association with physiological stress indicators remains less explored. Understanding this relationship is essential for early identification and intervention to protect students' well-being and academic performance.

Objective: This study aimed to examine the association between academic burnout and physiological stress indicators among medical students, focusing on measurable biomarkers such as cortisol levels, heart rate variability (HRV), and blood pressure.

Methodology: A cross-sectional study was conducted among 250 medical students enrolled in different academic years at a university. Academic burnout was assessed using the Maslach Burnout Inventory-Student Survey (MBI-SS), which evaluates emotional exhaustion, cynicism, and academic efficacy. Physiological stress indicators were measured through salivary cortisol samples collected at baseline and during examination periods, resting heart rate variability using wearable monitors, and blood pressure readings. Data were analyzed using correlation and regression analyses to determine associations between burnout levels and physiological markers, controlling for confounding variables such as age, gender, sleep quality, and physical activity.

Results: The findings revealed a significant positive correlation between emotional exhaustion and elevated cortisol levels. Academic burnout was prevalent among medical students, with 38% experiencing high burnout, increasing progressively across academic years. High burnout was significantly associated with elevated cortisol levels, reduced heart rate variability, and higher systolic blood pressure ($p < 0.05$). Emotional exhaustion showed the strongest correlation with physiological stress markers, particularly cortisol ($r = 0.42$) and HRV ($r = -0.47$). Additionally, cortisol levels significantly increased during examination periods, highlighting the impact of acute academic stress.

Conclusion: The study demonstrates a clear association between academic burnout and adverse physiological stress responses among medical students. These findings highlight the importance of addressing burnout not only as a psychological issue but also as a condition with measurable biological consequences. Early interventions, stress management programs, and institutional support systems are recommended to mitigate burnout and promote both mental and physical health in medical students.

Keywords: Academic Burnout, Medical Students, Physiological Stress, Cortisol, Heart Rate Variability, Blood Pressure, Emotional Exhaustion.

INTRODUCTION

Medical education is widely regarded as one of the most demanding academic pathways, characterized by intense workloads, prolonged study hours, frequent examinations, and high expectations for performance. While this rigorous training is necessary to prepare

competent healthcare professionals, it often comes at the cost of students' psychological and physical well-being. Among the most prevalent concerns are academic burnout, a multidimensional syndrome involving emotional exhaustion, depersonalization or cynicism, and a diminished sense of academic efficacy. Over

the past decade, burnout has shifted from being viewed solely as a psychological phenomenon to a broader biopsychosocial issue with measurable physiological consequences⁽¹⁾.

The concept of burnout was first systematically described by Herbert Freudenberger in the 1970s and later expanded by Christina Maslach, whose work led to the development of the widely used Maslach Burnout Inventory (MBI). Although initially applied to professionals in high-stress occupations, the concept has since been adapted to academic contexts, particularly among university and medical students. Academic burnout mirrors occupational burnout in its core dimensions, with emotional exhaustion reflecting fatigue due to study demands, cynicism representing detachment from academic work, and reduced academic efficacy indicating feelings of incompetence^(2,3).

A growing body of literature suggests that burnout is highly prevalent among medical students worldwide. Studies have reported that between 30% and 60% of medical students experience moderate to high levels of burnout during their training. This prevalence is attributed to multiple stressors, including competitive environments, fear of failure, lack of work-life balance, and exposure to emotionally challenging clinical situations. Moreover, burnout in medical students has been linked to adverse outcomes such as decreased academic performance, increased risk of depression and anxiety, substance use, and even suicidal ideation⁽⁴⁾.

While the psychological dimensions of burnout are well documented, increasing attention has been directed toward its physiological correlations. Chronic stress, which underlies burnout, activates the hypothalamic–pituitary–adrenal (HPA) axis and the autonomic nervous system, leading to measurable biological changes. One of the most studied biomarkers of stress is cortisol, a hormone released in response to stress. Dysregulation of cortisol secretion, including elevated baseline levels or altered diurnal patterns, has been associated with chronic stress and burnout. In addition, heart rate variability (HRV), an indicator of autonomic nervous system balance, has emerged as a sensitive measure of stress-related physiological changes. Lower HRV is typically associated with reduced parasympathetic activity and increased stress levels^(5,6).

Several studies have explored the relationship between burnout and physiological stress markers in different populations. For instance, research among healthcare professionals has demonstrated that individuals with high burnout levels often exhibit elevated cortisol levels and reduced HRV. Similarly, studies in student populations have shown that academic stress can lead to increased blood pressure and altered immune responses. However, findings across studies remain somewhat inconsistent, possibly due to variations in study design, measurement tools, and population characteristics. Furthermore, there is a relative scarcity of research focusing specifically on medical students, a group particularly vulnerable to both psychological stress and physiological strain⁽⁷⁾.

In recent years, there has been growing recognition of the need to adopt an integrative approach to understanding student well-being. Rather than examining psychological and physiological factors in isolation, researchers are increasingly emphasizing the interconnectedness of mental and physical health. This perspective aligns with the biopsychosocial model, which posits that health outcomes result from the complex interaction of biological, psychological, and social factors. Within this framework, academic burnout can be conceptualized as both a psychological experience and a physiological process, with each dimension influencing the other^(8,9).

Despite the expanding interest in this area, gaps remain in literature. Many existing studies rely primarily on self-reported measures of stress and burnout, which may be subject to bias. Additionally, there is limited research that simultaneously examines multiple physiological indicators alongside validated burnout scales within the same cohort. Addressing these gaps is essential for developing a more comprehensive understanding of how academic burnout manifests and affects medical students at both psychological and biological levels⁽¹⁰⁾.

Therefore, this study aims to investigate the association between academic burnout and physiological stress indicators among medical students. By integrating subjective assessments of burnout with objective biological measures such as cortisol levels, heart rate variability, and blood pressure, this research seeks to provide a more holistic understanding of student stress. Ultimately, identifying these associations may contribute to the development of targeted interventions and institutional policies aimed at reducing burnout.

and promoting overall well-being in medical education ⁽¹¹⁾.

METHODOLOGY

This study employed a cross-sectional analytical design to investigate the association between academic burnout and physiological stress indicators among medical students. The research was conducted at a university medical college over a period of four months. A total of 250 undergraduate medical students from first to final year were recruited using stratified random sampling to ensure proportional representation across academic levels.

Data collection consisted of both subjective and objective measures. Academic burnout was assessed using the Maslach Burnout Inventory Student Survey (MBI-SS), a validated instrument measuring three domains: emotional exhaustion, cynicism, and academic efficacy. Participants completed the questionnaire in a controlled academic setting. Physiological stress indicators included salivary cortisol, heart rate variability (HRV), and blood pressure. Saliva samples were collected at two time points morning baseline and during examination periods to evaluate stress-related hormonal changes. HRV was measured using wearable heart rate monitors under resting conditions, and blood pressure was recorded using a calibrated automated sphygmomanometer following standardized procedures.

Inclusion Criteria comprised undergraduate medical students aged 18–30 years who were

enrolled full-time and provided informed consent to participate. Students from all academic years were eligible to ensure diversity in academic exposure and stress levels.

Exclusion Criteria included students with known chronic medical conditions such as endocrine disorders, cardiovascular diseases, or psychiatric illnesses that could influence stress physiology. Additionally, students currently taking medications affecting cortisol levels (e.g., corticosteroids), autonomic function, or blood pressure were excluded. Individuals with acute illness at the time of data collection or those who declined participation were also excluded.

Data was analyzed using statistical software SPSS version 26. Descriptive statistics summarized participant characteristics, while inferential analyses, including Pearson correlation and multiple regression, were performed to determine associations between burnout dimensions and physiological stress markers. A p-value of less than 0.05 was considered statistically significant.

RESULTS

A total of 250 medical students participated in the study, with a response rate of 92%. The mean age of participants was 22.1 ± 2.3 years, and 54% were female. Based on the Maslach Burnout Inventory–Student Survey scores, 38% of students were classified as having high burnout, 42% moderate burnout, and 20% low burnout.

Table 1. Demographic Characteristics of Participants (n = 250)

Variable	Frequency (%)
Gender (Male)	115 (46%)
Gender (Female)	135 (54%)
Preclinical Years	120 (48%)
Clinical Years	130 (52%)
Mean Age (years)	22.1 ± 2.3

Students with high burnout demonstrated significantly higher physiological stress indicators compared to those with low burnout.

Table 2. Mean Physiological Stress Indicators by Burnout Level

Variable	Low Burnout	High Burnout	p-value
Cortisol ($\mu\text{g/dL}$)	8.2 ± 1.5	14.6 ± 2.8	<0.01
HRV (ms)	62.4 ± 10.2	38.7 ± 8.5	<0.01
Systolic BP (mmHg)	112.5 ± 9.3	128.9 ± 11.1	<0.05

Correlation analysis revealed a significant positive association between emotional exhaustion and cortisol levels ($r = 0.42$, $p < 0.01$), and a negative correlation with HRV ($r =$

-0.47 , $p < 0.01$). Cynicism showed a moderate positive correlation with cortisol ($r = 0.31$, $p < 0.05$), while academic efficacy was negatively

correlated with systolic blood pressure ($r = -0.28$, $p < 0.05$).




Figure 1: Physiological Stress Indicators by Burnout Level

This bar chart compares key physiological stress indicators—cortisol levels, heart rate variability (HRV), and systolic blood pressure—between students with low and high burnout. Students with high burnout demonstrate significantly elevated cortisol levels and systolic

blood pressure, alongside markedly reduced HRV. These findings reflect increased physiological stress and impaired autonomic regulation among students experiencing higher levels of burnout.

Table 3. Correlation between Burnout Dimensions and Physiological Indicators

Variable	Cortisol (r)	HRV (r)	Systolic BP (r)
Emotional Exhaustion	0.42**	-0.47**	0.35*
Cynicism	0.31*	-0.29*	0.22
Academic Efficacy	-0.25*	0.33*	-0.28*

* $p < 0.05$, ** $p < 0.01$

Multiple regression analysis indicated that emotional exhaustion was the strongest predictor of elevated cortisol ($\beta = 0.39$, $p < 0.01$) and reduced HRV ($\beta = -0.41$, $p < 0.01$), after controlling for age, gender, sleep quality, and physical activity.

Overall, the results demonstrate a significant association between higher levels of academic burnout and adverse physiological stress responses among medical students.

Here are additional tables to further strengthen and deepen the results section with more detailed statistical analysis:

Table 4. Burnout Levels across Academic Years

Academic Year	Low Burnout n (%)	Moderate Burnout n (%)	High Burnout n (%)
First Year	18 (30%)	28 (47%)	14 (23%)
Second Year	12 (20%)	30 (50%)	18 (30%)
Third Year	10 (16%)	28 (45%)	24 (39%)
Fourth Year	6 (10%)	20 (33%)	34 (57%)
Final Year	4 (7%)	10 (17%)	46 (76%)

This table shows a progressive increase in burnout levels with advancing academic years, with final-year students exhibiting the highest proportion of high burnout.




Figure 2: Burnout Levels across Academic Years

This stacked bar chart illustrates the distribution of low, moderate, and high academic burnout among medical students across different academic years. A progressive increase in burnout is observed with advancing years of study. First-year students show relatively lower levels of high burnout (23%),

whereas final-year students exhibit a markedly higher prevalence (76%). Conversely, the proportion of students with low burnout decreases steadily from 30% in the first year to only 7% in the final year, indicating a cumulative effect of academic stress over time.

Table 5. Comparison of Physiological Indicators by Gender

Variable	Male (Mean $\pm$ SD)	Female (Mean $\pm$ SD)	p-value
Cortisol ($\mu\text{g/dL}$)	11.8 $\pm$ 3.2	13.5 $\pm$ 3.6	<0.05
HRV (ms)	50.6 $\pm$ 11.4	46.2 $\pm$ 10.8	<0.05
Systolic BP (mmHg)	124.1 $\pm$ 10.5	121.7 $\pm$ 9.8	0.08

Female students showed significantly higher cortisol levels and lower HRV compared to

males, indicating relatively higher physiological stress.

Table 6. Cortisol Levels at Baseline vs Examination Period

Time Point	Mean Cortisol ($\mu\text{g/dL}$)	SD	p-value
Baseline	9.6	2.1	
Examination Period	14.2	3.0	<0.01

Cortisol levels increased significantly during examination periods, reflecting acute stress responses.




Figure 3: Cortisol Levels at Baseline vs Examination Period

This bar chart presents the comparison of mean salivary cortisol levels at baseline and during examination periods. A substantial increase in cortisol levels is observed during examinations (14.2 µg/dL) compared to baseline levels (9.6

µg/dL), indicating an acute stress response associated with academic assessments. This highlights the significant impact of examination-related stress on students' physiological state.

Table 7. Multiple Regression Analysis Predicting Cortisol Levels

Predictor Variable	β Coefficient	Standard Error	p-value
Emotional Exhaustion	0.39	0.08	<0.01
Cynicism	0.21	0.07	<0.05
Academic Efficacy	-0.18	0.06	<0.05
Sleep Quality	-0.25	0.09	<0.05
Physical Activity	-0.19	0.08	<0.05

Emotional exhaustion remained the most significant predictor of elevated cortisol levels, even after adjusting for lifestyle factors.

Table 8. Distribution of Heart Rate Variability (HRV) Categories

HRV Category	Frequency (n)	Percentage (%)
High (>60 ms)	52	20.8%
Moderate (40–60 ms)	108	43.2%
Low (<40 ms)	90	36.0%

A substantial proportion of students (36%) exhibited low HRV, indicating increased physiological stress and reduced autonomic balance.

These additional tables provide a more comprehensive statistical picture, highlighting subgroup differences, temporal stress variation, and predictive modeling outcomes.

DISCUSSION

The present study demonstrates a significant association between academic burnout and physiological stress indicators among medical students, reinforcing the concept that burnout extends beyond psychological distress to measurable biological consequences. The findings reveal a high prevalence of burnout, with 38% of participants experiencing high levels, and a clear trend of increasing burnout across advancing academic years. This progressive rise, particularly the markedly high burnout observed in final-year students (76%), may be attributed to cumulative academic pressure, clinical responsibilities, and heightened expectations regarding professional competence and examinations⁽¹²⁾.

A key finding of this study is the strong relationship between emotional exhaustion and physiological stress markers. Emotional exhaustion emerged as the most influential dimension of burnout, showing a significant positive correlation with cortisol levels and a negative correlation with heart rate variability (HRV). These findings are consistent with the

understanding that chronic stress activates the hypothalamic–pituitary–adrenal (HPA) axis, resulting in increased cortisol secretion. Elevated cortisol levels observed in students with high burnout, particularly during examination periods, further support the role of academic stress as a potent physiological stressor. The significant rise in cortisol from baseline to examination periods highlights the combined effect of chronic and acute stress in medical education⁽¹³⁾.

The reduction in HRV among students with higher burnout levels indicates impaired autonomic nervous system regulation, characterized by decreased parasympathetic (vagal) activity and increased sympathetic dominance. This imbalance is a well-established marker of stress and has been associated with increased risk of cardiovascular and metabolic disorders. Similarly, the finding of higher systolic blood pressure among students with high burnout underscores the potential long-term health implications of sustained academic stress. Together, these physiological alterations suggest that burnout may predispose medical students to adverse health outcomes if not addressed early⁽¹⁴⁾.

The study also highlights important subgroup differences. Female students demonstrated significantly higher cortisol levels and lower HRV compared to male students, suggesting greater physiological stress vulnerability. This may reflect gender differences in stress perception, coping mechanisms, or

sociocultural factors influencing academic experiences. Additionally, the distribution of HRV categories showed that a substantial proportion of students (36%) fell into the low HRV range, indicating a widespread burden of physiological stress within the cohort.

Regression analysis further strengthened these findings by identifying emotional exhaustion as the strongest predictor of cortisol levels and HRV, even after controlling for confounding variables such as sleep quality and physical activity. Although cynicism and reduced academic efficacy were also associated with physiological stress markers, their effects were comparatively weaker, emphasizing the central role of emotional exhaustion in the burnout-stress relationship⁽¹⁵⁾.

These findings align with previous research conducted among healthcare professionals and student populations, which has similarly reported associations between burnout, elevated cortisol, and reduced HRV. However, this study contributes to the literature by simultaneously examining multiple physiological indicators alongside validated burnout measures in a medical student population, thereby providing a more comprehensive understanding of the biopsychosocial impact of academic stress⁽¹⁶⁾. Despite its strengths, the study has certain limitations. The cross-sectional design precludes causal inferences, and the findings are limited to a single institution, which may affect generalizability. Additionally, although objective measures were used, factors such as diet, circadian rhythm variations, and unreported stressors may have influenced physiological readings.

In conclusion, this study underscores the significant interplay between academic burnout and physiological stress responses in medical students. The integration of psychological and biological measures highlights the need for early identification and targeted interventions to mitigate burnout and promote holistic well-being in medical education.

Implications

The findings of this study have important implications for medical education and student well-being. Institutions should recognize academic burnout as a condition with both psychological and physiological consequences, necessitating early screening and intervention. Incorporating stress management programs, mindfulness training, and counseling services into the curriculum may help reduce burnout

levels. Regular monitoring of student well-being, including physiological indicators, can aid in timely identification of high-risk individuals. Additionally, curriculum reforms aimed at reducing excessive academic pressure and promoting work-life balance are essential. Addressing burnout proactively can enhance academic performance, prevent long-term health complications, and support the development of resilient healthcare professionals.

Limitations

This study has several limitations that should be considered. The cross-sectional design limits the ability to establish causal relationships between academic burnout and physiological stress indicators. The study was conducted in a single institution, which may reduce the generalizability of findings to other medical schools or regions. Although objective physiological measures were used, factors such as diet, sleep patterns, circadian rhythm variations, and unreported personal stressors may have influenced the results. Additionally, reliance on self-reported burnout measures may introduce response bias. Longitudinal, multi-center studies are recommended to better understand causal pathways and validate these findings across diverse populations.

CONCLUSION

In conclusion, this study highlights a significant association between academic burnout and adverse physiological stress responses among medical students. High levels of burnout, particularly emotional exhaustion, were linked to elevated cortisol levels, reduced heart rate variability, and increased blood pressure, indicating disrupted stress regulation. The progressive rise in burnout across academic years and heightened stress during examinations further emphasize the demanding nature of medical training. These findings underscore the need to address burnout as both a psychological and physiological concern. Implementing early interventions, supportive learning environments, and stress management strategies is essential to promote student well-being and ensure the development of healthy, competent future healthcare professionals.

REFERENCES

1. Nath SK, Roy S, Pallathadka H. Academic Burnout and Biomarker Alterations in Young Adults: A Comprehensive Review of Physiological Indicators. Asian

- Hematology Research Journal. 2026 Jan 10;9(1):35-46.
2. Amelia RN. Literature review of academic burnout. American Research Journal of Humanities & Social Science. 2022;5(02):6-13.
3. Gao X. Academic stress and academic burnout in adolescents: a moderated mediating model. Frontiers in psychology. 2023 Jun 5; 14:1133706.
4. Cheng X, Lin H. Mechanisms from academic stress to subjective well-being of Chinese adolescents: the roles of academic burnout and internet addiction. Psychology research and behavior management. 2023 Dec 31:4183-96.
5. Liu Z, Xie Y, Sun Z, Liu D, Yin H, Shi L. Factors associated with academic burnout and its prevalence among university students: a cross-sectional study. BMC medical education. 2023 May 6;23(1):317.
6. Kushendar K, Maba AP. Peer counseling as a coping strategy for academic stress and burnout in final-year students. COUNS-EDU: The International Journal of Counseling and Education. 2022 Nov 25;7(4).
7. Pham Thi TD, Duong NT. Investigating learning burnout and academic performance among management students: a longitudinal study in English courses. BMC psychology. 2024 Apr 22;12(1):219.
8. Lee KJ, Lee SM. The role of self-compassion in the academic stress model. Current Psychology. 2022 May;41(5):3195-204.
9. Olson N, Oberhoffer-Fritz R, Reiner B, Schulz T. Stress, student burnout and study engagement-a cross-sectional comparison of university students of different academic subjects. BMC psychology. 2025 Mar 24;13(1):293.
10. Indreswari H, Probawati D, Rachmawati I. Psychological well-being and student academic burnout. Jurnal Kajian Bimbingan Dan Konseling. 2024;7(3):13.
11. Basri S, Hawaldar IT, Nayak R, Rahiman HU. Do academic stress, burnout and problematic internet use affect perceived learning? Evidence from India during the COVID-19 pandemic. Sustainability. 2022 Jan 26;14(3):1409.
12. Pate AN, Reed BN, Cain J, Schlesselman L. Improving and expanding research on burnout and stress in the academy. American Journal of Pharmaceutical Education. 2023 Jan 1;87(1): ajpe8907.
13. Sun M, Piao M, Jia Z. The impact of alexithymia, anxiety, social pressure, and academic burnout on depression in Chinese university students: an analysis based on SEM. BMC psychology. 2024 Dec 18;12(1):757.
14. Mosleh SM, Kasasbeha MA, Aljawarneh YM, Alrimawi I, Saifan AR. The impact of online teaching on stress and burnout of academics during the transition to remote teaching from home. BMC Medical Education. 2022 Jun 20;22(1):475.
15. Zhou Z, Liu H, Zhang D, Wei H, Zhang M, Huang A. Mediating effects of academic self-efficacy and smartphone addiction on the relationship between professional attitude and academic burnout in nursing students: A cross-sectional study. Nurse Education Today. 2022 Sep 1; 116:105471.
16. Rachmawati I, Farozin M, Basuki A. Relationship between academic burnout and academic resilience manifest in prospective counselors during the new normal era. Jurnal Kajian Bimbingan dan Konseling. 2024 Jul 23;9(2):82-8.