

Research Article

Comparative Study of Surgical Outcome of Different Approaches in 90 Patients of Varicocele at Gmc Baramulla, Jammu and Kashmir

Sheikh Mehmood Rashid¹, Owais Khan², Sheikh Aamir Rashid^{3*}

¹M.S. Associate Professor Department of General Surgery Govt. Medical College and Associated Hospital Baramulla, Kashmir (India) – 193101.

²Post Graduate Scholar, Department of General Surgery Govt. Medical College and Associated Hospital Baramulla, Kashmir (India) – 193101.

^{3*}Post Graduate Scholar Department of Psychiatry Govt. Medical College Srinagar, – 190001.

Correspondence Author: Dr. Shiekh Aamir Rashid

Post Graduate Scholar Department of Psychiatry Govt. Medical College Srinagar, – 190001.

Email: mahmoodshiekh234@gmail.com

Received: 16.03.26, Revised: 18.04.26, Accepted: 23.05.26

ABSTRACT

Background: Varicocele is the most commonly diagnosed prepubertal andrological condition and is an abnormal dilatation, lengthening and tortuosity of the veins of the spermatic cord. The clinical presentation of varicocele may vary from dull and dragging unilateral or bilateral testicular pain to visible varicose veins lying over the hemiscrotum.

Methods: A total number of 90 male adolescents with varicoceles were treated at the department of General Surgery at Govt. Medical College Baramulla Kashmir from January 2018 to December 2023 by three recognized surgical techniques viz., scrotal, trans inguinal and high ligation (Palomo's technique) and the results of surgical outcome in the three mentioned surgical approaches were compared.

Conclusion: We concluded that trans inguinal method is the best one for the correction of varicocele with easy approach, fewer complications and shorter hospital stay. There was a marked improvement in sperm quality and testicular volume in patients operated by this route.

Results: 56.6% of patients belonged to age group 15-20 years and in 90% of cases, left sided testis was involved. Among 6 patients who were infertile 2 (33.3%) patients became fertile and 4 (66.6%) patients had no effect on their fertility potential after varicocele correction. A significant change in testicular volume was noticed post operatively in all the 3 methods. A statistically significant improvement in sperm count (>50 million per ml) was observed in Trans inguinal technique while a significant improvement in sperm motility (>60%) was seen in trans inguinal and Palomo's approach.

Keywords: Varicocele, Testis, Sperm Count, Ligation, Surgical Approaches.

INTRODUCTION

Varicocele by definition means a compact pack of vessels filled with melancholic blood and is an abnormal dilatation, lengthening and tortuosity of the veins of the spermatic cord[1]. The reported incidence in the general male population ranges from 8 to 23% [2-5]. The veins ascend in three longitudinal groups, the anterior or spermatic group (varicocele), the middle or vassal group and the posterior or cremasteric group. As the veins ascend from the scrotum, they decrease in number, increase in size and enter the deep inguinal ring as the

internal spermatic veins [6-9]. Varicocele occurs on the left side in as many as 98% of affected patients [10-14]. There is a high degree of correlation between the grade of varicocele, effect on testicular volume and results following varicocelectomy[15-18].

The reported incidence of varicocele in infertile males is as high as 30% suggesting that varicocele might play a significant role in infertility [19]. This study involved an analysis of 30 patients treated by three modalities of treatment for varicoceles; viz, the scrotal, Trans

inguinal and Palomo's procedure and the results of treatment were analysed with reference to effects on semen analysis, testicular size, fertility outcome, pregnancy rates and complications.

MATERIAL AND METHODS

The study included 90 male adolescents with their varicocele who presented at department of General surgery, Govt. Medical College Baramulla, Kashmir from January 2018 to December 2023. Cases were analysed according to age of partner, duration of infertility, previous pregnancies, pregnancy outcome, and history of any previous systemic and genitourinary disorders. All patients were subjected to a complete physical examination. Pre-operatively haemogram, routine urine examination, urine culture, ESR, kidney function tests, sperm analysis and assessment of testicular size (testicular volume) by calipers and ultrasonography was done in all the study subjects. Semen volume, sperm concentration, percentage and progression of motility and morphology were studied and this preoperative semen analysis was compared with three postoperative semen samples at 3,6 and 9 months. The effects of treatment were also studied by measuring testicular volume which was compared with preoperative reading. The results of the three surgical techniques for the treatment of varicocele viz scrotal, Trans inguinal and Palomo's or high ligation were assessed by the recurrence rate complications and effects on semen analysis.

RESULTS AND DISCUSSION

In the present study of 90 patients with varicocele, maximum number of cases (56.6%) was in age group 15-20 years and in 81 (90%) cases left sided testis was involved. Of total number of 90 patients, 24 (26%) patients were married and out of them 18 (75%) patients were infertile.

Among 18 patients who were married but infertile, 6(33.3%) patients became fertile and 12 (66.6%) patients had no effect on their fertility potential after varicocele correction. Preoperative evaluation revealed 66 (73.3%) patients with sperm count of <50 million/ml

and 24 (226.6%) patients with sperm count of ≥ 50 million/ml; 69 (76.6%) patients with sperm motility of <60% motile sperms and 21 (23.3%) patients with sperm motility of. $\geq 60\%$ motile sperms and 63(70%) patients with sperm morphology showing <60% oval forms and 27(30%) patients with that of $\geq 60\%$ oval forms. Of 181 patients with left sided varicoceles testicular volume was <15 ml in 48 (59.25%) cases and ≥ 15 ml in 33 (40.74%) cases while in 9 patients with right sided varicoceles, 6 (66.6%) patients were with testicular volume of <15 ml and 3 (33.3%) case with that of ≥ 15 ml.

In our series three modalities of treatment were used for varicocele repair each group consisting of 30 (33.3%) patients.

Postoperative evaluation of semen analysis done at 3,6 and 9 months revealed the results of sperm count, sperm motility and sperm morphology as shown in tables 1, 2 and 3 respectively. The number of cases with sperm count of >50 million was statistically significant at 9 months. Also a statistically significant increase in number of cases with sperm motility of >60% at 9 months was seen. However, no significant improvement in sperm morphology (oval forms) was observed.

A significant change in testicular volume was noted postoperatively reaching statistical significance at 1% level. Two tailed P value (Fishers exact test) was >0.2 (insignificant). The results of postoperative semen analysis in the patients operated by the three mentioned recognized surgical methods for correction of varicocele were analysed and the results of sperm count, sperm motility and sperm morphology are shown in table 4,5 and 6 respectively. A statistically significant improvement ($P < 0.005$) in sperm count (>50 million/ml) was observed in trans inguinal technique only in sperm count (>50 million/ml) while a significant improvement in sperm motility (>60%) was seen in trans inguinal and Palomos approach. No significant improvement in sperm morphology occurred with either of the 3 techniques. Out of all the 3 surgical techniques for treatment of varicocele, maximum incidence of complications was seen

in scrotal method and minimum incidence in Trans inguinal method as shown in table 7.

DISCUSSION

In the prospective randomized controlled study of the three methods for the correction of varicocele in the 90 patients studied in the present series, sperm analysis showed a marked improvement after varicolectomy. Of the total number of 90 patients, 50% showed sperm count of ≥ 50 million/ ml ($p < .01$), 66.6% of the patients showed sperm motility of $\geq 60\%$ ($p < 0.002$) and 40% showed a morphology of $\geq 60\%$ ($p > 0.5$) oval forms. Our results are in similarity with work done by Arnold M balcer et al 1981, Dubin L et al 1970, Kass EJ et al 1987 and Haans LC et al 1991[20-23]. There was marked improvement in testicular volume postoperatively (77.7% of patients showed a testicular volume of ≥ 15 ml with $p = 0.01$). Our results are in accordance with those of Kass and Balman 1984, Pintus C et al 2001, Beddy Petal 2005, Orda R et al 1987, Reitelman C et al 1987, Fertil Steril 1985, Lec J et al 2008 and Dobanovacki D 2010[24-31].

Trans inguinal method was found to be the best one for correction of varicocele with fewer complications and shorter hospital stay. There was a marked improvement in sperm quality

and testicular volume postoperatively by this method. Similar results were supported by Vermeulen A et al 1984, Li F et al 2012 and Miyaoka R et al 2012[32-34].

Scrotal method should not be usually employed for correction of varicocele because of its major complications. There was no effect on semen analysis in patients operated by this route.

CONCLUSION

Varicocele is a common disorder that affects adolescent population. Its clinical presentation can be widely variable and often asymptomatic or silent. If left untreated, varicocele can cause testicular damage leading to infertility. Thus, urgent clinical attention should be sought to correct and repair varicocele. Various surgical procedures have been proposed but trans inguinal method was found to be the best one for correction of varicocele with less complications and shorter hospital stay and marked improvement in sperm quality and testicular volume by this method. Scrotal method has been found to be associated with major complications and should be avoided.

- Sponsorship and Financial support – Nil
- Conflicts of Interest – None
- Written Consent of study subjects taken and there are no ethical issues.

Table -1 Sperm Count (Preoperative and Postoperative Results) in 90 Patients

Sperm Count	Preoperative	Postoperative		
	No (%)	At 3 mon. Follow-up no. (%)	At 6 mon. Follow-up no. (%)	At 9 mon. Follow-up no. (%)
<50 million	66 (73.3)	60 (66.6)	51 (56.6%)	36 (40%)
>50 million	24 (26.6)	30 (33.3)	39 (43.3%)	54 (60%)
Yates corrected (χ^2)		.08	1.17	5.50
Df		1	1	1
p		1S	1S	Sig

1S = Insignificant, Sig = Significant

Table-2 Sperm Motility (Preoperative and Postoperative Results)

Sperm Count	Preoperative	Postoperative		
	No. (%)	At 3 mon. Follow-up no. (%)	At 6 mon. Follow-up no. (%)	At 9 mon. Follow-up no. (%)
<60% motile	69 (73.3)	54 (60)	42 (46.6)	30 (33.3)
$\geq 60\%$ motile	21 (23.3)	36 (40)	48 (53.3)	60 (66.6)

Yates corrected (χ^2)		1.233	4.51	9.7
Df		1	1	1
p		>0.2	>0.3	<.002
		1S	1S	HS

1S = Insignificant, HS = Highly Significant

<60%	63 (70)	69 (76.6)	63 (70)	54 (60)
>60%	27 (30)	21 (23.3)	27 (30)	36 (40)
Yates corrected (χ^2)		0.09	0.08	0.29
Df		1	1	1
p		>0.2	>0.3	<.002
		1S	1S	1S

1S = Insignificant

Table-4 Sperm Count (Results of the Three Surgical Methods)

		Pre-operative	Post-Operative	P value
Method	No. of patients	Sperm count (million/ml) <50 >50 no.(%) no.(%)	Sperm count (million/ml) <50 >50 no.(%) no.(%)	(2- tailed fisher's Exact Test)
Trans inguinal	30	24(80) 06(20)	03(10) 09(90)	<0.005(Sig)
Scrotal	30	21(70) 09(30)	21(70) 03(30)	>0.9(1S)
Palomo's	30	21(70) 09(30)	09(30) 07(70)	>0.1(1S)

1S = Insignificant, Sig = Significant

Table-5 Sperm Motility (Results of the Three Surgical Methods)

		Pre-operative	Post-Operative	P value
Method	No. of patients	Sperm count (million/ml) <50 >50 no.(%) no.(%)	Sperm count (million/ml) <50 >50 no.(%) no.(%)	(2- tailed fisher's Exact Test)
Trans inguinal	30	27(90) 01(10)	03(10) 09(90)	<0.001(Sig)
Scrotal	30	18(60) 04(40)	18(60) 04(40)	>0.09(1S)
Palomo's	30	24(80) 02(20)	03(10) 09(90)	>0.003(Sig)

1S = Insignificant, Sig = Significant

Table-6 Sperm Morphology-Oval Forms (Results of the Three Surgical Methods)

Method	No. of Patients	Pre-Operative	Post-Operative	P value (2-Tailed Fisher's Exact Test)
---------------	------------------------	----------------------	-----------------------	---

		Sperm Morphology <60% >/60% no.(%) no.(%)	Sperm Morphology <60% >/60% no.(%) no.(%)	
Trans Inguinal	30	15(30) 05(50)	09(30) 07(70)	>0.4(1S)
Scrotal	30	15(50) 05(50)	15(50) 05(50)	>0.9(1S)
Palomo's	30	15(40) 06(60)	09(30) 07(70)	>0.5(1S)

1S = Insignificant

Table-7 Complications in Relation to the Method Used

Table-3 Sperm Morphology (Preoperative and Postoperative)						Meth od	No. of Patie nts	Hydr oce le no.(%)	Scrot al haem ato ma no.(%)	Wou nd infe ctio n no.(%)	Varico cele Recurr renc e no.(%)	Varic oce le Persis ten ce no.(%)
Results)												
Spe rm	Preoper ative	Postoperative (oval forms)										
Cou nt	(oval forms)											
	No. (%)	At 3 mon ths	At 6 mon ths	At 9 mo nths								
		Follo w- up no. (%)	Foll ow- up no.(%)	Foll ow- up no. (%)								
		Tran s Ingui nal	30	-	-	03(1 0)	-	03(10)				
		Scrot al	30	09(3 0)	09(3 0)	03(1 0)	03(10)	06(20)				

Palomo's	30	03(10)	-	06(20)	03(10)	06(20)
----------	----	--------	---	--------	--------	--------

REFERENCES

1. Ham AW. Histology, Lippincott, Philadelphia Montreal 1965; pp 964-65.
2. Palomo A. Radical cure of varicocele by anew technique; preliminary report. J. Urol 1949; 61: :604-7.
3. Rivilla F, Casillas AJ. G, Gallego J et al. Percutaneous venography and embolization of the internal spermatic vein by spring coil for treatment of the left varicocele in children. Journal of paediatric Surgery April, 1995; 30(4)
4. Pryor J. L and Howards S. S. Varicocele, Urologic Clinics of North America August, 1987;
5. Risser W. I., Lipshultz I.1. Frequency of varicocele in black adolescents. J. Adolesc Health Care January, 1981; 5(1); 28-29.
6. Etriby AA, Ibrahim AS, Mahoud K et al. Subfertility and varicocele: Venogram demonstration of anastomosis sites in subfertile men. Steril 1975; 26: 1013-17.
7. Gill B.K.J. Stanley, Maldonado J, Reda E and Levitt B,. S. Significance of intra-operative venographic patterns on the post-operative recurrence and surgical incision placement of paediatric varicoceles, Journal of Urology August, 1990; 144: 502-5.
8. Glezerman M, Rakowszezyk M, Lunenfeld B et al. Varicocele in oligospermic patients pathophysiology and results after ligation and dissection

- of the internal spermatic vein. *Journal of Urology* May, 1976; 115.
9. Gohary M.A. Boyhood varicocele. *Br. J. Hosp. Med.* March, 1986; 35(30); 183-5.
10. Dubin L and Amelar R.D. Varicocele size and results of varicocelectomy in selected subfertile men with varicocele. *Fertil, Steril* 1970; 21: 606-9.
11. Coolsact B. IR.A. The varicocele syndrome: Venography determining the optimal level for surgical management. *Journal of Urology* Dec. 1980; 124.
12. Burington R.S. (cited by Joel Safyan 1988). *Handbook of mathematical tables and formulas*, 5th edition New York McGraw-Hill 1973;21.
13. Bloodgood J.C. (quoted by Richard Szabo and Robert Kessler 1984). A warning against operations for varicocele on applicants for enlistment, registrants for the selective draft and soldiers *J.A.M.A* 1918; 70:409.
14. Berger O.G. Varicocele in adolescence. *Clin Pediatr (Phila)* Dec., 1980; 19(12):810-11.
15. McClue R.D.Robaire B. Relationship of varicocele size to testicular volume and to effect on varicocelectomy outcome. Presented at annual meeting of American urological Association, Las Vegas (abstract 504)1983.
16. Sathya Sriniv, Belur Veerachari S. Does varicocelectomy improve gonadal function in men with hypogonadism and infertility? Analysis of a prospective study. *Int J Endocrinol* 2011;2011:916380.
17. Shiraishi K, Matsuyama H, Takihara H. Pathophysiology of varicocele in male fertility in the era of assisted reproductive technology. *Int J Urol* 2012;19:538-50.
18. Laven JS, Haans LC, Mali WP, te Velde ER, Wensing CJ, Eimers JM. Effects of varicocele treatment in adolescents, a randomized study. *Fertil Steril* 1992;58:756-62.
19. Dubin L, Amelar RD. Varicocele. *Urol Clin North Am* 1978;5:563-72.
20. Balkar A.M. The varicocele and male fertility. *Urologic Clinics of North America* 1981;8(1).
21. Dubin L, Amelar RD. Varicocele size and results of varicocelectomy in selected subfertile men with varicocele. *Fertil Steril* 1970;21:606-9.
22. Kass EJ, Belman AB. Reversal of testicular growth failure by varicocele ligation. *J Urol* 1987;137:475-6.
23. Haans LC, Laven JS, Mali WP, te Velde ER Wensing CJ, Testis volumes, semen quality, and hormonal patterns in adolescents with and without a varicocele. *Fertil Steril* 1991;56:731-6.
24. Kass E.J. and Belman AB. Reversal of testicular growth failure by varicocele ligation. *Journal of Urology* March, 1987;137.
25. Pintus C, Rodriguez Matas MJ, ManzooniC, Nanni L, Perrelli L. Varicocele in pediatric patients: Comparative assessment of different surgical approaches. *Urology* 2001;57:154-7.
26. Beddy P, Geoghegan T, Browne RF, Torreggiani WC. Testicular varicoceles. *Clin Radiol* 2005;60:1248-55.
27. Orda R, Safyan J, Manor H, Witz E, Sofer Y. Diagnosis of varicocele and postoperative evaluation using inguinal ultrasonography. *Ann Surg* 1987;206:99-101.
28. Reitelman C, Burbique K. A., Sawezuk I.S. et al Diagnosis and surgical correction of the paediatric varicocele. *Journal of Urology* October, 1987; 138.
29. Comparison among different methods for the diagnosis of varicocele. World Health Organization. *Fertil Steril* 1985;43:575-82.
30. Lee J, Binsaleh S, Lo K, Jarvi K. Varicoceles: The diagnostic dilemma. *J Androl* 2008;29:143-6.
31. Dobanovacki D. Varicocele in adolescents. *Med Pregl* 2010;63:741-6.
32. Vermeulen A, Vandeweghe M. Improved fertility after varicocele correction: Fact or fiction. *Fer Sterl* August, 1984; 42(2).

33. LiF, Yue H, Yamaguchi K, Okada K, Matsushita K, Ando M, et al. Effect of surgical reoair on testosterone production in infertile men with varicocele: A meta-analysis. *Int J Urol* 2012;19:149-54.
34. Miyaoka R, Esteves SC. A critical appraisal on the role of varicocele in male infertility. *Adv Urol* 2012;2012:597495.