

Research Article

A PROSPECTIVE STUDY OF EPIDEMIOLOGY, CAUSES AND PROGNOSIS OF ACUTE RENAL FAILURE IN A TERTIARY CARE HOSPITAL

Dr Lingaraju K S¹, Dr. Pavan Kumar D S², Dr Deepak D^{3*}

¹Assistant Professor, Department of General Medicine, Subbaiah Institute of Medical Sciences, Holehonnur road, Purle.

²MBBS MD DNB General Medicine, Subbaiah Institute of Medical Sciences, Shivamogga.

^{3*}Assistant Professor, General Medicine, Subbaiah Institute of Medical Sciences, Shimoga.

Corresponding Author: Dr Deepak D

Assistant Professor, General Medicine, Subbaiah Institute of Medical Sciences, Shimoga.

Received date: 10-07-2026, Date of Acceptance: 15-07-2026, Date of Publication: 18-07-2026.

Abstract

Introduction: AKI, previously known as Acute Renal Failure, is a clinical syndrome characterized by an abrupt decline in glomerular filtration rate sufficient to decrease the elimination of nitrogenous waste products (urea and creatinine) and other uremic toxins. Acute kidney injury (AKI) is a common clinical syndrome with a broad range of etiological profile. It is associated with major morbidity and significant mortality due to the severity of the causative illness. This study is to determine the various causes of AKI, in our hospital and to find out the incidence of AKI by using renal failure indices and to analyze outcome of AKI pertaining to the aetiology.

Materials and Methods: Study is conducted on 200 consecutive patients admitted in Department of General Medicine, Subbaiah Institute of Medical Sciences, Holehonnur road, purle From March 2024 to February 2025. All AKI patients aged between 15 to 70 years on hemodialysis were included in the study. Persons with CKD were excluded from

study. In all these patients, detailed clinical history and detailed clinical examination was carried out. Co-morbid conditions such as diabetes mellitus (DM), hypertension (HTN); Baseline and peak levels of serum creatinine, urine output was documented. Data regarding laboratory investigations were collected to confirm the etiology of AKI, which included complete urine analysis, metabolic panel, hematologic profile. Coagulation profile and blood culture (wherever necessary). Radiological tests, ultrasonography, urinalysis or urine culture, Requirement of RRT was noted.

Results: The age distribution of these patients ranged from 15 years to 70 years. The highest number of cases in age group 51 to 60 year (32%). Average age is 56 ± 5.6 years. The mean age among men was 55.22 ± 8.72 years and in women was 56.45 ± 4.61 years. The male to female ratio was 1.3:1. Fever, nausea and oliguria are the most common clinical features seen 66, 66 and 64 percentages respectively, followed by edema (23%) and loose stools (17%).

Conclusion: AKI remains a common disorder among critically ill patients Consistent with other studies from developing world; this study has also shown that infections, nephrotoxins and gastroenteritis are the primary causes of AKI at our institute. Most of these causes can be prevented with simple interventions such as health education on oral rehydration, quality prenatal and emergency obstetric care, appropriate management of infections and taking appropriate precautions when prescribing potentially nephrotoxic medications.

Key Words: Acute kidney Injury, diabetes mellitus, hypertension, gastroenteritis.

INTRODUCTION

AKI, previously known as Acute Renal Failure, is a clinical syndrome characterized by an abrupt decline in glomerular filtration rate sufficient to decrease the elimination of nitrogenous waste products (urea and creatinine) and other uremic toxins.¹

Acute kidney injury (AKI) is a common clinical syndrome with a broad range of etiological profile. It is associated with major morbidity and significant mortality due to the severity of the causative illness. This study is to determine the various causes of AKI, in our hospital and to find out the incidence of AKI by using renal failure indices and to analyze outcome of AKI pertaining to the aetiology.²

Acute kidney injury (AKI) affects one in five hospitalized patients. The epidemiology of AKI in developing countries is unique in that certain causes, such as the infections, obstetric causes and nephrotoxins, which are largely obsolete in developed countries remain important causes.³

In India, acute kidney injury constitutes 1.5% of all general hospital admissions, of which 60% are due to medical causes. The most common causes of AKI are: acute diarrheal diseases, sepsis, infection (malaria, UTI, pneumonia, viral hepatitis), snake bite, cardiac failure, diabetes mellitus, nephrotoxic drug use, malignancy, SLE, hypertension etc. major surgery like exploratory laparotomy, Whipple's procedure etc. are also an important cause of AKI.⁴ Advanced age, liver diseases, underlying comorbid illness (DM, HTN, IHD, COPD, cirrhosis) have been implicated as risk factor for the development of AKI. AKI constitutes approximately 5% of hospital admissions and up to 30% of admissions to intensive care units. Detection of the incidence, etiological profile and outcome of AKI is important for commencement of preventive and therapeutic strategies, geographical, etiological, cultural and economic variations determine the dissimilarities among patterns of AKI in different regions of the world. Although reliable statistics on the prevalence of AKI are not available, statistics on referrals to dialysis units suggest that the condition is more common in India as compared to the West. There is an increase in the incidence of surgical and sepsis-related AKI. Hence we conducted a prospective study to look for pattern of AKI in our locality.⁵

MATERIALS AND METHODS

Study is conducted on 200 consecutive patients admitted in Department of General Medicine, Subbaiah Institute of Medical Sciences, Holehonnur road, purle From March 2024 to February 2025.

Inclusion criteria

All AKI patients aged between 15 to 70 years on hemodialysis.

Exclusion criteria

Persons with CKD were excluded from study.

200 consecutive patients admitted in Department of General Medicine, Subbaiah Institute of Medical Sciences, Holehonnur road, purle with AKI All patients aged above 15 years with features of AKI as per AKIN (Acute Kidney Injury Network) criteria. Which is defined as an increase in serum creatinine of 0.3 mg/dl or more within 48 hours of observation were taken into

study after obtaining written informed consent. In all these patients, detailed clinical history and detailed clinical examination was carried out. Co-morbid conditions such as diabetes mellitus (DM), hypertension (HTN); Baseline and peak levels of serum creatinine, urine output was documented. Data regarding laboratory investigations were collected to confirm the etiology of AKI, which included complete urine analysis, metabolic panel, hematologic profile. Coagulation profile and blood culture (wherever necessary). Radiological tests, ultrasonography, urinalysis or urine culture, Requirement of RRT was noted.

RESULTS

The age distribution of these patients ranged from 15 years to 70 years. The highest number of cases in age group 51 to 60 year (32%). Average age is 56 ± 5.6 years. The mean age among men was 55.22 ± 8.72 years and in women was 56.45 ± 4.61 years. The male to female ratio was 1.3:1.

Gender	No of patients	Percentage
Male	130	65
Female	70	35

Table 1: Gender distribution

Age group	No of patients	Percentage
15-20	4	2
21-30	10	5
31-40	24	12
41-50	54	27
51-60	64	32
61-70	44	22

Table 2: Age distribution

Clinical features	No. of patients	Percentage
Oliguria	128	64
Loose stools	34	17
Vomiting	132	66
Fever	132	66
Burning micturition	12	6
Jaundice	8	4
Dyspnoea	20	10
Altered Sensorium	10	5
Edema	46	23

Table 3: Clinical presentations

Fever, nausea and oliguria are the most common clinical features seen 66, 66 and 64 percentage respectively, followed by edema (23%) and loose stools (17%).

Diseases	No. of patients	Percentage
Diabetes mellitus	66	33
Hypertension	56	35
COPD	10	5
Coronary artery disease	16	8

Table 4: Co-morbid conditions

Test	Value	Percentage
Urea (mg/dl)	<100	56
	>100	44
Creatinine(mg/dl)	<4	54
	>4	46
WBC count	<12000	35
	>12000	65
Potassium (mmol/l)	<5	58
	>5	42
Hb (g/dl)	<10	54
	>10	46

Table 5: Biochemical tests.

Etiology	Subgroup	Number of cases	Total
Medical	Sepsis	50	150
	Gastroenteritis	24	
	Cardiogenic/CCF	16	
	CVA	6	
	Chronic liver disease	12	
	Malaria	4	
	Nephrotoxic medications	14	
	Paraquet poisoning	12	

Table 6: Causes for AKI.

DISCUSSION

There were 130 males (65%) and 70 females (35%) in the present study. The male to female ratio was 1.3:1. This finding is consistent with that of Olaga B et al 60% males and 40% females with a ratio of 1.6:1 and in study by Kapadia MP et al. It is 73% males and 27% females.⁶

The age distribution of these patients ranged from 15 years to 70 years with 2/3rd being in between 41 to 70 years. The maximum number of cases in age group 51 to 60 year i.e 32%. Average age is 56 ± 5.6 years. This is with Mataloun et al mean age is 55.3 years and In Picard study it is 59.5 years. In Ibrahim A et al study mean age is 36.7 ± 14.5 years.⁷

Clinical features are comparable with other studies particularly in relation to fever and oliguria. Incidence of sepsis and gastroenteritis were comparable to many other Indian studies.

A large proportion of medical causes in our study are due to a high incidence of infections, especially gastrointestinal diseases, malaria with multi organ dysfunction. The other conditions were snake bite, paraquet poisoning and chronic liver diseases. The obstetrics and surgical causes are similar to other studies.⁸

Medical causes contributed most to AKI it is 75%, surgical causes in 14% and obstetric causes in 11% of patients. In Umesh L et al it is 73.7%, 6.4% and 19.8% respectively. Low incidence of surgical causes in our setup can be attributed to non-performance of cardiac surgery and transplant surgery in our institute.⁹

Only 20% of our patients required hemodialysis for recovery. Remaining were

treated conservatively. Mortality was around 25% mainly due to co morbid conditions sepsis, cardiac, liver and CNS disorder rather than due to renal failure itself. It is comparable to study done by Kashinkunti MD et al which is 28.3%. In Umesh L et al study it is 9.61%.¹⁰

ARF in the setting of ICU is characterized by increasing mortality, high incidence sepsis, multi-organ failure and mortality. Sepsis and multi-organ failure are the major causes of mortality in these patients. These findings alarm for early detection and aggressive management of sepsis and its associated complications so as to bring down the mortality in patients admitted to intensive care unit. It is important for preventive measures such as restoration of intravascular volume status, maintenance of adequate renal perfusion pressures, and limiting exposure to nephrotoxins.

CONCLUSION

AKI remains a common disorder among critically ill patients Consistent with other studies from developing world; this study has also shown that infections, nephrotoxins and gastroenteritis are the primary causes of AKI at our institute. Most of these causes can be prevented with simple interventions such as health education on oral rehydration, quality prenatal and emergency obstetric care, appropriate management of infections and taking appropriate precautions when prescribing potentially nephrotoxic medications. Future studies may also benefit by better identifying modifiable risk factors to prevent the development of AKI from the outset. The timely and aggressive management will certainly reduce the incidence of AKI.

REFERENCES

1. ChughKS. Changing trends in acute renal failure in third-world countries-Chandigarh study. *Q J Med.* 1989;73(272):1117-23.
2. Thadani R, Pascaul M, Bonvette VB. Acute renal failure. *New EngJ Med.* 1996;30;1448-51.
3. Liu KD, Chertow GM. Acute renal failure. In: Fauci AS, Braunwald E, Kaspar DL, Hauser SL, Longo DL, Jameson JL, editors. *Harrison's Principles of Internal Medicine.* 17th ed. New York: McGraw Hill; 2008:1752-1761.
4. Kaufman J, Dhakal M, Patel B, Hamburger J. Community-acquired acute renal failure. *Am J Kidney Dis.* 1991;17:191-8.
5. Jayakumar M, Prabakar MR, Fernando EM, Manojan R, Venkatraman R, Balaraman V. Epidemiologic trend changes in acute renal failure-a tertiary center experience from South India. *Ren Fail.* 2006;28:405-10.
6. Anandh U, Renuka S, Somiah S, Vincent L. Acute renal failure in the tropics: emerging trends from a tertiary care hospital in South India. *ClinNephrol.* 2003;59:341-4.
7. Molitoris BA, Levin A, Warnock DG. Improving outcomes from acute kidney injury. *J Am SocNephrol.* 2007;18:1992-4.
8. Balafoutis O, Andrikos E, Tseke P, Tsintas A, Pappas E, Kokkolou E, et al. Outcome and epidemiology of hospital-acquired acute renal failure. *Bantao J.* 2007;5(2):55-7.
9. Kapadia MP. A study of clinical profile of patients with acute kidney injury in a tertiary care centre. *Int J Adv Scient Res.* 2016;2(8):160-6.
10. Mataloun SE, Machado FR, Senna APR, Guimarães HP, Amaral JLG. Incidence, risk factors and prognostic factors of acute renal failure in patients admitted to an intensive care unit; *Crit Care.* 2003;7(Suppl 3):45.